

**Just In Time Quick Check**  
**Standard of Learning (SOL) A.7d**

**Strand: Functions**

**Standard of Learning (SOL) A.7d**

*The student will investigate and analyze linear and quadratic function families and their characteristics both algebraically and graphically, including intercepts.*

**Grade Level Skills:**

- Identify the domain, range, zeros, and intercepts of a function presented algebraically or graphically.
- Use the x-intercepts from the graphical representation of a quadratic function to determine and confirm its factors.
- Investigate and analyze characteristics and multiple representations of functions with a graphing utility.

**Just in Time Quick Check**

**Just in Time Quick Check Teacher Notes**

**Supporting Resources:**

- VDOE Mathematics Instructional Plans (MIPS)
  - [A.7bcd - Functions 2: Exploring Quadratic Functions](#) (Word) / [PDF Version](#)
  - [A.7cd - Quadratic Connections](#) (Word) / [PDF Version](#)
  - [A.7cd - Solving Linear Equations Using Functions with Desmos](#) (Word) / [PDF Version](#)
- VDOE Algebra Readiness Formative Assessments
  - [A.7c,d](#) (Word) / [PDF](#)
- VDOE Word Wall Cards: Algebra I ([Word](#)) | ([PDF](#))
  - x-Intercepts
  - Parent Functions - Linear, Quadratic
- VDOE Rich Mathematical Tasks: The Soccer Competition
  - [A.7 The Soccer Competition Task Template](#) (Word) / [PDF Version](#)
- Desmos Activities
  - [Transforming Lines](#)
  - [Two Truths and a Lie: Quadratics](#)
  - [What's my Transformation?](#)
  - [Polygraph: Parabolas, Polygraph: Parabolas Part 2](#)
  - [Polygraph: Quadratics](#)
  - [Will It Hit the Hoop?](#)

**Supporting and Prerequisite SOL:** [A.1b](#), [A.4a](#), [A.6c](#), [8.16b](#), [8.16d](#), [8.17](#), [7.10c](#), [7.12](#)

### SOL A.7d - Just in Time Quick Check

1. Circle all of the following functions that have an  $x$ -intercept of 3.

$$f(x) = x^2 - 2x + 3$$

$$g(x) = 2x - 6$$

$$h(x) = x^2 - 9$$


$$p(x) = -\frac{1}{2}x + 3$$

$$q(x) = -3x^2 + 10x - 3$$

2. Let  $g(x) = -\frac{2}{3}x + 5$  and  $h(x) = \frac{4}{5}x + k$ . For which value of  $k$  will the  $x$ -intercept of  $g(x)$  be equivalent to the  $x$ -intercept of  $h(x)$ ?

3. Write the  $x$ - and  $y$ -intercept of the function  $f(x) = 3x - 4$  each as an ordered pair.

4. Circle the  $y$ -intercept of the function shown on the graph.



5. Which of the following functions have exactly one  $x$ -intercept?

$$f(x) = 4x(x - 5)$$

$$g(x) = x^2 - 6x + 9$$

$$h(x) = 2x^2 + 4x + 3$$

$$j(x) = -3(x + 1)$$

## SOL A.7d - Just in Time Quick Check Teacher Notes

### Common Errors/Misconceptions and their Possible Indications

1. Circle all of the following functions that have an x-intercept of 3.

$$f(x) = x^2 - 2x + 3$$

$$g(x) = 2x - 6$$

$$h(x) = x^2 - 9$$

$$p(x) = -\frac{1}{2}x + 3$$

$$q(x) = -3x^2 + 10x - 3$$

*A common error a student may make is to select the functions with a y-intercept of 3, such as  $f(x)$  and  $p(x)$ . This may indicate that a student has difficulty differentiating between an x-intercept and y-intercept using an algebraic approach. A strategy that might be useful is to have a student represent the functions visually and determine which functions have an x-intercept of 3 then make the connection algebraically. Desmos is a powerful tool that can be used to show connections between algebraic forms, graphs, and intercepts.*


2. Let  $g(x) = -\frac{2}{3}x + 5$  and  $h(x) = \frac{4}{5}x + k$ . For which value of  $k$  will the x-intercept of  $g(x)$  be equivalent to the x-intercept of  $h(x)$ ?

*A common error that a student may make is to say that  $k = 7.5$ , which is the x-intercept of  $g(x)$ . This indicates the student would benefit from additional practice in comparing functions and working with constant variable terms. A strategy that could be used is for the students to experiment with the slider feature in Desmos to demonstrate what happens to the graph and equation of  $h(x)$  as  $k$  changes in value.*

3. Write the x- and y-intercept of the function  $f(x) = 3x - 4$  each as an ordered pair.

*A common error a student may make is to write the x-intercept as  $(0, \frac{4}{3})$  or the y-intercept as  $(-4, 0)$ . This may indicate a misunderstanding of representing x- and y-intercepts as ordered pairs. A strategy that might be helpful for students is to verify the intercepts using a graphing utility such as Desmos. In addition, a student might find helpful to use the table feature in Desmos to verify intercepts.*

4. Circle the y-intercept of the function shown on the graph.



*A common error a student may make is to circle both the  $x$ - and  $y$ -intercepts or to circle only the  $x$ -intercepts. This may indicate that a student has difficulty distinguishing between  $x$ - and  $y$ -intercepts and a misunderstanding between intercepts and solutions of a function. A strategy that might be helpful for students is to represent the  $x$ - and  $y$ -intercepts as a set of ordered pairs or as a table to show the similarities and differences between the coordinates.*

5. Which of the following functions have exactly one  $x$ -intercept?

$$f(x) = 4x(x - 5)$$

$$g(x) = x^2 - 6x + 9$$

$$h(x) = 2x^2 + 4x + 3$$

$$j(x) = -3(x + 1)$$

*A common error a student may make is to select  $f(x)$  as having only one  $x$ -intercept because it is written in factored form and appears to have one binomial factor. This may indicate that a student does not recognize that the GCF of  $4x$  is also a factor of the function and constitutes a unique  $x$ -intercept. A strategy that might be helpful for students is to verify the intercepts using a graphing utility such as Desmos. In addition, a student might find helpful to use the table feature in Desmos to verify the  $x$ -intercepts a function.*